



TITLE:

SPECTRAL THEORY FOR DIVERGENCE-FORM OPERATORS (Spectral and Scattering Theory and Related Topics)

AUTHOR(S):

BEN-ARTZI, MATANIA

CITATION:

BEN-ARTZI, MATANIA. SPECTRAL THEORY FOR DIVERGENCE-FORM OPERATORS (Spectral and Scattering Theory and Related Topics). 数理解析研究所講究録 2008, 1607: 77-84

ISSUE DATE:

2008-07

URL:

<http://hdl.handle.net/2433/139969>

RIGHT:

SPECTRAL THEORY FOR DIVERGENCE-FORM OPERATORS

MATANIA BEN-ARTZI

Workshop on "Spectral and Scattering Theory and Related Topics"
Research Institute for Mathematical Sciences, Kyoto University, Japan, January 2008

1. Introduction and Statement of Results

Let $H = - \sum_{i,j=1}^n \partial_i a_{i,j}(x) \partial_j$, where $a_{i,j} = a_{j,i}$, be a formally self-adjoint operator in $L^2(\mathbb{R}^n)$, $n \geq 2$, where the notation $\partial_j = \frac{\partial}{\partial x_j}$ has been used.

We assume that the real measurable matrix function $a(x) = \{a_{i,j}(x)\}_{1 \leq i,j \leq n}$ satisfies, with some positive constants $a_1 > a_0 > 0$, $\Lambda_0 > 0$,

$$(1.1) \quad a_0 I \leq a(x) \leq a_1 I, \quad x \in \mathbb{R}^n,$$

$$(1.2) \quad a(x) = I \quad \text{for } |x| > \Lambda_0.$$

In what follows we shall use the notation $H = -\nabla \cdot a(x) \nabla$.

We retain the notation H for the self-adjoint (Friedrichs) extension associated with the form $(a(x) \nabla \varphi, \nabla \psi)$, where (\cdot, \cdot) is the scalar product in $L^2(\mathbb{R}^n)$. When $a(x) \equiv I$ we get $H = H_0 = -\Delta$.

Let

$$R_0(z) = (H_0 - z)^{-1}, R(z) = (H - z)^{-1}, \quad z \in \mathbb{C}^\pm = \{z/ \quad \pm \operatorname{Im} z > 0\},$$

be the associated resolvent operators.

The purpose of this paper is to study the continuity properties of $R(z)$ in certain operator topologies, as z approaches the real axis. To fix the ideas, we shall generally assume that $\operatorname{Im} z > 0$, with obvious modifications for $\operatorname{Im} z < 0$.

Definition 1.1. Let $[\alpha, \beta] \subseteq \mathbb{R}$. We say that H satisfies the "Limiting Absorption Principle" (LAP) in $[\alpha, \beta]$ if $R(z)$, $z \in \mathbb{C}^+$, can be extended continuously to $\operatorname{Im} z = 0$, $\operatorname{Re} z \in [\alpha, \beta]$, in a suitable operator topology. In this case we denote the limiting values by $R^+(\lambda)$, $\alpha \leq \lambda \leq \beta$.

A similar definition applies for $z \in \mathbb{C}^-$, but the limiting values $R^-(\lambda)$ will be, generally speaking, different from $R^+(\lambda)$. Observe that the precise specification of the operator topology in the above definition is left open. Typically, it will be the uniform operator topology associated with weighted- L^2 or Sobolev spaces, which will be introduced later.

It is well-known that our assumptions (1.1), (1.2) imply that $\sigma(H)$, the spectrum of H , is the half-axis $[0, \infty)$, and is entirely absolutely continuous. The "threshold" $z = 0$ plays a special role in this setting, as we shall see later. Thus, consider first the case $[\alpha, \beta] \subseteq (0, \infty)$.

Under assumptions close to ours here (but also assuming that $a(x)$ is continuously differentiable) a weaker version (roughly, "strong" instead of "uniform" convergence of the resolvents) was obtained by Eidus [14, Theorem 4 and Remark 1]. For $H = H_0$ the LAP has been established by Agmon [1]. Indeed, it was established for operators of the type $H_0 + V$, where V is a short-range perturbation. However, an inspection of Agmon's perturbation-theoretic proof shows that it cannot be extended to our operator H , in a straightforward way. Observe on the other hand that the short-range potential V can be replaced by a potential depending only on direction $(x/|x|)$ [15] or a perturbation of such a potential [23, 24]. In this case the condition $\alpha > 0$ is replaced by $\alpha > \limsup_{|x| \rightarrow \infty} V(x)$. The LAP for the periodic

case (namely, $a(x)$ is symmetric and periodic) has recently been established in [22]

Note that in this case the spectrum is absolutely continuous and consists of a union of intervals ("bands").

We also refer to [16] where the existence and completeness of the wave operators $W_{\pm}(H, H_0)$ is established under suitable smoothness assumptions on $a(x)$ (however, $a(x) - I$ is not assumed to be compactly supported and H can include also magnetic and electric potentials). Note that by a well-known theorem of Kato and Kuroda [19], if H, H_0 satisfy the LAP in $[\alpha, \beta]$ (with respect to the same operator topologies) then the wave operators over this interval exist and are complete.

In this paper we focus on the study of the LAP for H in $[\alpha, \beta]$ where $\alpha < 0 < \beta$. This case has been studied for the Laplacian H_0 [6, Appendix A] and in the one-dimensional case ($n = 1$) in [3, 4, 10]. The present paper deals with the multi-dimensional case $n \geq 2$.

Throughout this paper we shall make use of the following weighted- L^2 and Sobolev spaces. First, for $s \in \mathbb{R}$ and m a nonnegative integer we define.

$$(1.3) \quad L^{2,s}(\mathbb{R}^n) := \{u(x) \quad / \quad \|u\|_{0,s}^2 = \int_{\mathbb{R}^n} (1 + |x|^2)^s |u(x)|^2 dx < \infty\}$$

$$(1.4) \quad H^{m,s}(\mathbb{R}^n) := \{u(x) \quad / \quad D^{\alpha}u \in L^{2,s}, \quad |\alpha| \leq m, \quad \|u\|_{m,s}^2 = \sum_{|\alpha| \leq m} \|D^{\alpha}u\|_{0,s}^2\}$$

(we write $\|u\|_0 = \|u\|_{0,0}$).

More generally, for any $\sigma \in \mathbb{R}$, let $H^{\sigma} \equiv H^{\sigma,0}$ be the Sobolev space of order σ , namely,

$$(1.5) \quad H^{\sigma} = \{\hat{u} \quad / \quad u \in L^{2,\sigma}, \quad \|\hat{u}\|_{\sigma,0} = \|u\|_{0,\sigma}\}$$

where the Fourier transform is defined as usual by

$$\hat{u}(\xi) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} u(x) \exp(-i\xi x) dx.$$

For negative indices we denote by $\{H^{-m,s}, \|\cdot\|_{-m,s}\}$ the dual space of $H^{m,-s}$. In particular, observe that any function $f \in H^{-1,s}$ can be represented (not uniquely) as

$$(1.6) \quad f = f_0 + \sum_{k=1}^n i^{-1} \frac{\partial}{\partial x_k} f_k, \quad f_k \in L^{2,s}, \quad 0 \leq k \leq n.$$

In the case $n = 2$ and $s > 1$, we define

$$L_0^{2,s}(\mathbb{R}^2) = \{u \in L^{2,s}(\mathbb{R}^2) \mid \hat{u}(0) = 0\},$$

and set $H_0^{-1,s}(\mathbb{R}^2)$ to be the space of functions $f \in H^{-1,s}(\mathbb{R}^2)$ which have a representation (1.6) where $f_k \in L_0^{2,s}$, $k = 0, 1, 2$.

For any two normed spaces X, Y , we denote by $B(X, Y)$ the space of bounded linear operators from X to Y , equipped with the operator-norm topology.

The fundamental result obtained in the present paper is given in the following theorem.

THEOREM A. *Suppose that $a(x)$ satisfies (1.1), (1.2). Then the operator H satisfies the LAP in \mathbb{R} . More precisely, using the density of $L^{2,s}$ in $H^{-1,s}$, consider the resolvent $R(z) = (H - z)^{-1}$, $\text{Im } z \neq 0$, as two operator-valued functions, defined respectively in the lower and upper half-planes,*

$$(1.7) \quad z \rightarrow R(z) \in B(H^{-1,s}(\mathbb{R}^n), H^{1,-s}(\mathbb{R}^n)), \quad s > 1, \quad \pm \text{Im } z > 0.$$

Then these functions can be extended continuously from $\mathcal{C}^\pm = \{z \mid \pm \text{Im } z > 0\}$ to $\overline{\mathcal{C}^\pm} = \mathcal{C}^\pm \cup \mathbb{R}$, with respect to the operator-norm topology. In the case $n = 2$ replace $H^{-1,s}$ by $H_0^{-1,s}$.

In particular, it follows that the limiting values $R^\pm(\lambda)$ are continuous at $\lambda = 0$ and H has no resonance there. The study of the resolvent near the threshold $\lambda = 0$ is sometimes referred to as "low energy estimates". As mentioned earlier, this result has been established in the case $H = H_0$ [6, Appendix A]. The paper [25] deals with the two-dimensional ($n = 2$) case, but the resolvent $R(z)$ is restricted to continuous compactly supported functions f , thus enabling the use of pointwise decay estimates of $R(z)f$ at infinity. The case of the closely related "acoustic propagator", where the matrix $a(x) = b(x_1)I$ is scalar and dependent on a single coordinate, has been extensively studied [4, 9, 12, 17, 18, 20], as well as the "anisotropic" case where $b(x_1)$ is a general positive matrix [5]. The proof of the theorem will be given in Section 3. It is based on an extended version of the LAP for H_0 , with the resolvent $R_0(z)$ acting on elements of $H^{-1,s}$, for suitable positive values of s (see Section 2).

An important application of the LAP in the case of perturbations of the Laplacian is the derivation of an "eigenfunction expansion theorem", where the eigenfunctions are perturbations of plane waves $\exp(i\xi x)$ [1, 29]. We can use the LAP result of Theorem A in order to derive a similar expansion for the operator H . In fact, our generalized eigenfunctions are given by the following definition.

Definition 1.2. *For every $\xi \in \mathbb{R}^n$ let*

$$(1.8) \quad \begin{aligned} \psi_\pm(x, \xi) &= -R^\mp(|\xi|^2)((H - |\xi|^2) \exp(i\xi x)) = \\ &R^\mp(|\xi|^2)(\sum_{l,j=1}^n \partial_l(a_{l,j}(x) - \delta_{l,j})\partial_j) \exp(i\xi x). \end{aligned}$$

The generalized eigenfunctions of H are defined by

$$(1.9) \quad \varphi_\pm(x, \xi) = \exp(i\xi x) + \psi_\pm(x, \xi).$$

Remark 1.3. *We label the eigenfunctions as "generalized" because they do not belong to the Hilbert space $L^2(\mathbb{R}^n)$.*

In analogy with the eigenfunction expansion theorem for short or long range perturbations of the Laplacian [1, 29] we can now state an eigenfunction expansion theorem for the operator H . We assume $n \geq 3$ in order to simplify the statement of the theorem. As we show below (see Proposition ??) the generalized eigenfunctions are (at least) continuous in x , so that the integral in the statement makes sense.

THEOREM B

Suppose that $n \geq 3$ and that $a(x)$ satisfies (1.1), (1.2). For any compactly supported $f \in L^2(\mathbb{R}^n)$ define

$$(1.10) \quad (\mathbb{F}_\pm f)(\xi) = (2\pi)^{-\frac{n}{2}} \int_{\mathbb{R}^n} f(x) \overline{\varphi_\pm(x, \xi)} dx, \quad \xi \in \mathbb{R}^n.$$

Then the transformations \mathbb{F}_\pm can be extended as unitary transformations (for which we retain the same notation) of $L^2(\mathbb{R}^n)$ onto itself. Furthermore, these transformations "diagonalize" H in the following sense.

$f \in L^2(\mathbb{R}^n)$ is in the domain $D(H)$ if and only if $|\xi|^2(\mathbb{F}_\pm f)(\xi) \in L^2(\mathbb{R}^n)$ and

$$(1.11) \quad H = \mathbb{F}_\pm^* M_{|\xi|^2} \mathbb{F}_\pm,$$

where $M_{|\xi|^2}$ is the multiplication operator by $|\xi|^2$.

As is well-known from the theory of Schrödinger operators, the LAP and the eigenfunction expansion theorem provide powerful tools for the treatment of a wide array of related problems. Here we give one such application, dealing with global space-time estimates for a generalized wave equation.

We consider the equation

$$(1.12) \quad \frac{\partial^2 u}{\partial t^2} = Hu = - \sum_{i,j=1}^n \partial_i a_{i,j}(x) \partial_j u,$$

subject to the initial data

$$(1.13) \quad u(x, 0) = u_0(x), \quad \partial_t u(x, 0) = v_0(x), \quad x \in \mathbb{R}^n.$$

We next replace the assumptions (1.1), (1.2) by stronger ones as follows.

Let $g(x) = (g_{i,j}(x))_{1 \leq i,j \leq n}$ be a smooth Riemannian metric on \mathbb{R}^n such that

$$(1.14) \quad g(x) = I \quad \text{for } |x| > \Lambda_0.$$

and assume that

$$(1.15) \quad a(x) = g^{-1}(x) = (g^{i,j}(x))_{1 \leq i,j \leq n}.$$

We have the following theorem.

THEOREM C

Suppose that $n \geq 3$ and that $a(x)$ satisfies (1.14), (1.15). Assume further that the geometry defined by the metric g has no "trapped geodesics" [27]. Then for any $s > 1$ there exists a constant $C = C(s, n) > 0$ such that the solution to (1.12), (1.13) satisfies

$$(1.16) \quad \int_{\mathbb{R}} \int_{\mathbb{R}^n} (1 + |x|^2)^{-s} |u(x, t)|^2 dx dt \leq C[\|u_0\|_0^2 + \| |D|^{-1} v_0 \|_0^2],$$

where as usual $|D|^{-1}$ denotes multiplication by the symbol $|\xi|^{-1}$.

This estimate generalizes similar estimates obtained for the classical ($g = I$) wave equation [2, 21].

We do not provide proofs of the theorems, but we include below the treatment of the *unperturbed* operator H_0 . This treatment is already new in the sense that it extends the treatment of the LAP beyond the L^2 setting (see the statement of Theorem A).

2. The Operator $H_0 = -\Delta$

Let $\{E_0(\lambda)\}$ be the spectral family associated with H_0 , so that

$$(2.1) \quad (E_0(\lambda)h, h) = \int_{|\xi|^2 \leq \lambda} |\hat{h}|^2 d\xi, \quad \lambda \geq 0, \quad h \in L^2(\mathbb{R}^n).$$

Following the methodology of [7, 13] we see that the weak derivative $A_0(\lambda) = \frac{d}{d\lambda} E_0(\lambda)$ exists in $B(L^{2,s}, L^{2,-s})$ for any $s > \frac{1}{2}$ and $\lambda > 0$. (Here and below we write $L^{2,s}$ for $L^{2,s}(\mathbb{R}^n)$). Furthermore,

$$(2.2) \quad \langle A_0(\lambda)h, h \rangle = (2\sqrt{\lambda})^{-1} \int_{|\xi|^2 = \lambda} |\hat{h}|^2 d\tau,$$

where \langle, \rangle is the $(L^{2,-s}, L^{2,s})$ pairing and $d\tau$ is the Lebesgue surface measure. Recall that by the standard trace lemma we have

$$(2.3) \quad \int_{|\xi|^2 = \lambda} |\hat{h}|^2 d\tau \leq C \|\hat{h}\|_{H^s}^2, \quad s > \frac{1}{2}.$$

However, we can refine this estimate near $\lambda = 0$ as follows.

Proposition 2.1. *Let $\frac{1}{2} < s < \frac{3}{2}$, $h \in L^{2,s}$. For $n = 2$ assume further that $s > 1$ and $h \in L_0^{2,s}$. Then*

$$(2.4) \quad \int_{|\xi|^2 = \lambda} |\hat{h}|^2 d\tau \leq C \min(\lambda^\gamma, 1) \|\hat{h}\|_{H^s}^2,$$

where

$$(2.5) \quad 0 < \gamma < s - \frac{1}{2},$$

and $C = C(s, \gamma, n)$. (Actually we can take $\gamma = s - \frac{1}{2}$ if $s \leq 1$ and $n \geq 3$).

Proof. If $n \geq 3$, the proof follows from [8, Appendix]. If $n = 2$ and $1 < s < \frac{3}{2}$ we have, for $h \in L_0^{2,s}$,

$$|\hat{h}(\xi)| = |\hat{h}(\xi) - \hat{h}(0)| \leq C_{s,\delta} |\xi|^\delta \|\hat{h}\|_{H^s},$$

for any $0 < \delta < \min(1, s - 1)$. Using this estimate in the integral in the right-hand side of (2.4) the claim follows also in this case. \square

Combining Equations (2.2), (2.3) and (2.4) we conclude that,

$$(2.6) \quad \begin{aligned} | \langle A_0(\lambda)f, g \rangle | &\leq \langle A_0(\lambda)f, f \rangle^{\frac{1}{2}} \langle A_0(\lambda)g, g \rangle^{\frac{1}{2}} \\ &\leq C \min(\lambda^{-\frac{1}{2}}, \lambda^\eta) \|f\|_{0,s} \|g\|_{0,\sigma}, \quad f \in L^{2,s}, \quad g \in L^{2,\sigma}, \end{aligned}$$

where either

$$(i) \quad n \geq 3, \quad \frac{1}{2} < s, \sigma < \frac{3}{2}, \quad s + \sigma > 2 \quad \text{and} \quad 0 < 2\eta < s + \sigma - 2,$$

$$(2.7) \quad \text{or}$$

$$(ii) \quad n = 2, \quad 1 < s < \frac{3}{2}, \quad \frac{1}{2} < \sigma < \frac{3}{2}, \quad s + \sigma > 2, \quad 0 < 2\eta < s + \sigma - 2$$

$$\text{and} \quad \hat{f}(0) = 0.$$

In both cases, $A_0(\lambda)$ is Hölder continuous and vanishes at $0, \infty$, so as in [7] we obtain

Proposition 2.2. *The operator-valued function*

$$(2.8) \quad z \rightarrow R_0(z) \in \begin{cases} B(L^{2,s}, L^{2,-\sigma}), & n \geq 3, \\ B(L_0^{2,s}, L^{2,-\sigma}), & n = 2, \end{cases}$$

where s, σ satisfy (2.7), can be extended continuously from \mathcal{C}^\pm to $\overline{\mathcal{C}^\pm}$, in the respective uniform operator topologies.

We shall now extend this proposition to more general function spaces. Let $g \in H^{1,\sigma}$, where s, σ satisfy (2.7). Let $f \in H^{-1,s}$ have a representation of the form (1.6). Equation (2.2) can be extended in an obvious way to yield

$$(2.9) \quad i^{-1} \langle A_0(\lambda) \frac{\partial}{\partial x_k} f_k, g \rangle = (2\sqrt{\lambda})^{-1} \int_{|\xi|^2=\lambda} \xi_k \hat{f}_k(\xi) \overline{\hat{g}(\xi)} d\tau, \quad k = 1, \dots, n.$$

We therefore obtain

Proposition 2.3. *The operator-valued function of Proposition 2.2 is well-defined (and analytic) for nonreal z in the following functional setting.*

$$(2.10) \quad z \rightarrow R_0(z) \in \begin{cases} B(H^{-1,s}, H^{1,-\sigma}), & n \geq 3, \\ B(H_0^{-1,s}, H^{1,-\sigma}), & n = 2, \end{cases}$$

where s, σ satisfy (2.7). Furthermore, it can be extended continuously from \mathcal{C}^\pm to $\overline{\mathcal{C}^\pm}$, in the respective uniform operator topologies.

Proof. In view of (2.9) and the considerations preceding Proposition 2.2, since $g \in H^{1,\sigma}$, we have instead of (2.6),

$$(2.11) \quad \begin{aligned} | \langle A_0(\lambda) \frac{\partial}{\partial x_k} f_k, g \rangle | \\ \leq C \min(\lambda^{-\frac{1}{2}}, \lambda^\eta) \|f\|_{-1,s} \|g\|_{1,\sigma}, \quad f \in H^{-1,s}, \quad g \in H^{1,\sigma}, \end{aligned}$$

so that the claim holds true if $H^{1,-\sigma}$ is replaced by $H^{-1,-\sigma}$. However, using that $H_0 R_0(z) = I + z R_0(z)$ we see that also $H_0 R_0(z)$ can be extended continuously (as z approaches the real line from either half-plane) with values in $H^{-1,-\sigma}$. The conclusion of the proposition follows since the norm of $H^{1,-\sigma}$ is equivalent to the graph-norm of H_0 in $H^{-1,-\sigma}$. \square

REFERENCES

- [1] S. Agmon, *Spectral properties of Schrödinger operators and scattering theory*, Ann. Scuola Norm. Sup. Pisa, **2**(1975), 151–218.
- [2] M. Ben-Artzi, *Regularity and smoothing for some equations of evolutions* in "Nonlinear Partial Differential Equations and Their Applications; Collège de France Seminar", Longman Scientific, vol. XI, Eds. H. Brezis and J.L.Lions, (1994), 1-12.
- [3] M. Ben-Artzi, *On spectral properties of the acoustic propagator in a layered band*, J. Diff. Eqs. **136** (1997), 115-135.
- [4] M. Ben-Artzi, Y. Dermenjian and J.-C. Guillot, *Analyticity properties and estimates of resolvent kernels near thresholds*, Comm. Partial Diff. Eqs. **25** (2000), 1753-1770.
- [5] M. Ben-Artzi, Y. Dermenjian and A.Monsef, *Resolvent kernel estimates near thresholds*, Diff. Integr. Eqs. **19** (2006), 1-14.
- [6] M.Ben-Artzi and A. Devinatz, *The limiting absorption principle for a sum of tensor products applications to the spectral theory of differential operators*, J. d'Analyse Math. **43** (1983/84), 215-250.
- [7] M.Ben-Artzi and A. Devinatz, "The limiting absorption principle for partial differential operators", Memoirs of the AMS **66** (1987).
- [8] M. Ben-Artzi and S. Klainerman, *Decay and regularity for the Schrödinger equation*, J. d'Analyse Math. **58** (1992), 25-37.
- [9] A. Boutet de Monvel-Berthier and D. Manda, *Spectral and scattering theory for wave propagation in perturbed stratified media*, J. Math. Anal. Appl. **191** (1995), 137-167.
- [10] A. Cohen and T. Kappeler, *Scattering and inverse scattering for steplike potentials in the Schrödinger equation*, Indiana Univ. Math. J. **34** (1985), 127-180.
- [11] E. Croc and Y. Dermenjian, *Analyse Spectrale d'une Bande Acoustique Multistratifiée. Partie I: Principe d'absorption limite pour une stratification simple*, SIAM J. Math. Anal. **26** (1995), 880-924.
- [12] S. DeBièvre and W. Pravica, *Spectral analysis for optical fibers and stratified fluids I: The limiting absorption principle*, J. Func. Anal. **98** (1991), 404-436.
- [13] V.G. Deich, E.L. Korotayev and D.R. Yafaev, *Theory of potential scattering, taking into account spatial anisotropy*, J. of Soviet Math. **34** (1986), 2040-2050.
- [14] D. M. Eidus, *The principle of limiting absorption* in "American Mathematical Society Translations", Series 2, vol. **47**, Providence, 1965, 157-192. (original in Russian: Mat. Sb. **57**, (1962), pp. 13-44).
- [15] I. Herbst, *Spectral and scattering theory for Schrödinger operators with potentials independent of $|x|$* , Amer. J. Math. **113** (1991), 509-565.
- [16] T. Ikebe and T. Tayoshi, *Wave and scattering operators for second-order elliptic operators in \mathbb{R}^n* , Publ. RIMS Kyoto Univ. Ser. A **4** (1968), 483-496.
- [17] M. Kadowaki, *Low and high energy resolvent estimates for wave propagation in stratified media and their applications*, J. Diff. Eqs. **179** (2002), 246-277.
- [18] M. Kadowaki, *Resolvent estimates and scattering states for dissipative systems*, Publ. RIMS, Kyoto Univ. **38** (2002), 191-209.
- [19] T. Kato and S.T. Kuroda, *The abstract theory of scattering*, Rocky Mountain J. Math. **1** (1971), 127-171.
- [20] K. Kikuchi and H. Tamura, *The limiting amplitude principle for acoustic propagators in perturbed stratified fluids*, J. Diff. Eqs. **93** (1991), 260-282.
- [21] K. Mochizuki, *Scattering theory for wave equations with dissipative terms*, Publ. RIMS, Kyoto Univ. **12** (1976), 383-390.
- [22] M. Murata and T. Tsuchida, *Asymptotics of Green functions and the limiting absorption principle for elliptic operators with periodic coefficients*, J. Math. Kyoto Univ. **46** (2006), 713-754.
- [23] B. Perthame and L. Vega, *Morrey-Campanato estimates for Helmholtz equations*, J. Func. Anal. **164** (1999), 340-355.
- [24] B. Perthame and L. Vega, *Energy decay and Sommerfeld condition for Helmholtz equation with variable index at infinity*, Preprint (2002).
- [25] A.G.Ramm, *Justification of the limiting absorption principle in \mathbb{R}^2* , in "Operator Theory and Applications", Fields Institute Communications, vol. **25**, Eds. A.G. Ramm, P.N. Shivakumar, A.V. Strauss, AMS (2000), 433-440.

- [26] D. Robert, *Relative time-delay for perturbations of elliptic operators and semiclassical asymptotics*, J. Func. Anal. **126** (1994), 36-82.
- [27] D. Robert, *Asymptotique de la phase de diffusion a haute énergie pour des perturbations du second ordre du laplacien*, Ann. Scient. ENS, 4 serie, **25** (1992), 107-134.
- [28] D. Robert, *Relative time delay and trace formula for long range perturbations of Laplace operator*, in "Operator Calculus and Spectral Theory", Eds. M. Demuth, B. Gramsch and B.-W. Schulze, Birkhauser, "Operator Theory: Advances and Applications", Vol. 57 (1992), 245-253.
- [29] Y. Saito, "Spectral Representations for Schrödinger Operators with Long-Range Potentials", Lecture Notes in Mathematics, vol. 727, Springer-Verlag 1979.
- [30] G. Vodev, *Uniform weighted estimates of the resolvent of the Laplace-Beltrami operator on unbounded Riemannian manifolds*,

INSTITUTE OF MATHEMATICS, HEBREW UNIVERSITY, JERUSALEM 91904, ISRAEL
E-mail address: mbartzi@math.huji.ac.il